Case: 1:17-md-02804-DAP Doc #: 3836-6 Filed: 08/12/21 1 of 8. PageID #: 516862

EXHIBIT E



TEXTS: Now references are online
- No specific texts - use online
resources

School

School

School

School

PHARM. D'

Pharmaceutics

Pharmaceutics

Pharmaceutics

Pharmaceutics

Pharmaceutics

Pharmaceutics

Pharm. D'

Doctor's Deg. in Pharm.

Pharm. D'

Doctor's Deg. in Pharm.

Pharmaceutics

Pharmaceutics

Pharm. D'

Doctor's Deg. in Pharm.

Pharm. D'

Doctor's Deg. in Pharm.

Pharm. D'

Doctor's Deg. in Pharm.

Pharm. D'

Also Deal a general practice

Pharmaceutics

Pharmaceutics

Pharmaceutics

Pharm. D'

Pharm. D'

Doctor's Deg. in Pharm.

Pharm. D'

Doctor's Deg. in Pharm.

Pharmaceutics

Pharmaceutics

Pharmaceutics

Pharmaceutics

Pharm. D'

Doctor's Deg. in Pharm.

Pharm. D'

Pharm. D'

Pharm. D'

Doctor's Deg. in Pharm.

Pharm. D'

Pharm.



DeMay Exhibit


1

3/2/21 Carrie Campbell, RDR











DOCTORS
GOOD WORKING

PECATORISHIPS
W/ AREA DOCTORS

PAIN Manual doe's write

Majority

Syed Akhter - Zaidi - Pain Mat

Syed Akhter - Zaidi - no longer

Placticing

Samuel Nigro - No memory


Dean PAHR - Pain MANAg mt


Spoke w of disgnosis rode

of dosoge change
of add. med. for

Suragry recovery)







Oxycodone	Redacted - Confidential PH	The Total Action of the Position of the Positi	No.
Content Con	Care	The Total Action of the Position of the Positi	No.
Oxycodone	Oxycodone Hydromorphone Methadone Other (optional - district specific) Check boxes that apply to assist the pharmacist in determining if the prescription should be filled. Attach checklist to hard copy of fix. Valid government photo ID copied and attached to hard copy of fix. For effix, attach copy at pick- up. ID is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. Prescription must not be dispensed. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the prescription specific for the same medication as the prescription specific for the same medication for safe prescription specific for sa	The Total Action of the Position of the Positi	No.
Check boses that apply to assist the pharmacist is determining if the prescription should be filled. Attach checklist to hard copy of Rx. For eRx, attach copy at pick-up. ID is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. Prescription must not be dispensed. Patient has received this prescription from Walgreens before. This prescription is from the same prescriber for the same medication as the previous fill. The prescription is from the same prescriber for the same medication as the previous fill. The prescription is from the same prescriber for the same medication as the previous fill. The prescription is from the same prescriber for the same medication as the previous fill. The prescription is from the same prescriber for the same medication as the previous fill. The prescription is from the same prescriber for the same medication as the previous fill. The prescription is from the same prescriber for the same medication as the previous fill. The prescription is from the same prescriber for the same medication as the previous fill. The prescription is from the same prescriber for the same medication as the previous fill. The prescription above: Technician Initials Steps for Pharmacist to Complete Yes in the prescription is being filled on time. If your state the interpretation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances are regulates early refills of controlled substance prescriptions, follow your state's regulations. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or prescriptions ≥ 50 Morphine Milligram E	Check boxes that apply to assist the pharmacist in determining if the prescription should be filled. Attach checklist to hard copy of Rx. Valid government photo ID copied and attached to hard copy of Rx. For eRx, attach copy at pick- up. ID is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. Patient has received this prescription from Walgreens before. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is billed (If cash or a cash discount card, use caution). Not Net C. A PAD AN Patient does not appear intoxicated or under the influence of illicit drugs. Teatient does not appear intoxicated or under the influence of illicit drugs. Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient Talking Points #10-16 Refer to Opiold Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opiolds for Safer Dos If in your professional judgment a call to the prescriber's clinical staff is war	The Total Action of the Position of the Positi	No.
Valid government photo ID copied and attached to hard copy of Rx. For eRx, attach copy at pickup. ID is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, 2 prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (if cash or a cash discount card, use caution). Not Net. A 640 Patient does not appear intoxicated or under the influence of illicit drugs. 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes Note that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes Note that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes Note that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes Note that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes Note that I have used the Target Drug Good Faith Dispensing to Complete the Internal Patient Incomplete	Valid government photo ID copied and attached to hard copy of Rx. For eRx, attach copy at pickup. ID is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. Patient has received this prescription from Walgreens before. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validatest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validatest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validatest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validatest that I have used the Target Drug Good Faith Dispensing the checklist validation procedures to validatest that I have used the Target Drug Good Faith Dispensing to Checklist validation procedures to validatest that I have used the Target Drug Good Faith Dispensing ore met. Jeps Patient Amount of Sond Faith Dispensing ore met. Jeps Patient Checklist Validation procedures to validate the prescription without verification provided the of Good Faith Dispensing ore met. Jeps Patient Checklist Validation Provided the of Good Faith Dispensing ore met. Jeps Patient Validation Provided the of Good Faith Dispensing ore met. Jeps Patient Validation Provided the Of Good Faith Dispensing ore met. Jeps Patient Validation Provided the Of Good Faith Dispensing ore met. Jeps Patient Validation Provided the Of Good Faith Dispensing ore met. Jeps Patient Validation Provided the Of Control Patient Validation Provided the O	The Total Action of the Position of the Positi	No.
up. 10 is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. Patient has received this prescription from Walgreens before. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is billed (if cash or a cash discount card, use caution). Not Net. A PAD AND AND AND AND AND AND AND AND AND A	up. ID is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. Patient has received this prescription from Walgreens before. This prescription is from the same prescriber for the same medication as the previous fill. 3 and Party Insurance is billed (if cash or a cash discount card, use caution). Not Net. A peop at test that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opiold Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opiolds for Safer Dos if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	The state stat	No.
1 to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, 2 prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (if cash or a cash discount card, use caution). Not Net. A pub of the patient does not appear intoxicated or under the influence of illicit drugs. 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes M Tegulates early refills of controlled substance prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient Talking Points #10-16 Refer to Opiold Dosage Calculator and CDC handout "Calculating Total Dally Dose of Opiolds for Safer Dosage" If no vour professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing ore met.)	1 to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). Not Net. A PAD PATE of Patient does not appear intoxicated or under the influence of illicit drugs. 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	Test the Yes	
No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). Not Net. A 24D Medical of Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes Note that a validation in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances and be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opiold Dosage Calculator and CDC handout "Calculating Total Dally Dose of Opiolds for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). Not Net. A paper of Patient does not appear intoxicated or under the influence of illicit drugs. 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or glagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient Talking Points #10-16 Refer to Opiold Dosage Calculator and CDC handout "Calculating Total Dally Dose of Opiolds for Safer Dos if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	Test the Yes	
2 prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). Not Net. A PLD Mode of Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes Note that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes Note that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes Note that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes Note that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Use In Power Steps In Power	2 prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). Not Net. A PLO AN 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient Talking Points #10-16 Refer to Oploid Dosage Calculator and CDC handout "Calculating Total Dally Dose of Oploids for Safer Dos if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section, if no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	Test the Yes	
3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). Not Not. A PLO Mode of Patient does not appear intoxicated or under the influence of illicit drugs. 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the important on above: Technician Initials Steps for Pharmacist to Complete Yes Note that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the important of the prescription is being filled on time. If your state the pregulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances are not be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or glagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opiold Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opiolds for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section, if no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (if cash or a cash discount card, use caution). Not Net. A PAD Add Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Oploid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Oploids for Safer Dos if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	TO STATE THE TANK	No.
This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (if cash or a cash discount card, use caution). Not Net. A pub Mode of Patient does not appear intoxicated or under the influence of illicit drugs. 1 attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes Note that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the important of the prescription in itials to Complete Yes Note that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the important of the prescription is being filled on time. If your state that I have used the Target Drug Good Faith Dispensing medical provided the prescription is being filled on time. If your state the I have a patient of the prescription your state's regulations. Patient and/or prescripter address is within geographical proximity to pharmacy; any variances are not reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or glagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient of Talking Points #10-16 Refer to Opiold Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opiolds for Safer Dosage" if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (if cash or a cash discount card, use caution). Not Net. A PAD AN 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Oploid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Oploids for Safer Dos if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	TO STATE THE TANK	No.
Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≠ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Oploid Dosage Calculator and CDC handout "Calculating Total Dally Dose of Oploids for Safer Dosage" if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	5 3rd Party Insurance is billed (if cash or a cash discount card, use caution). Not Net. A PAD AM 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dos if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	Yes	No Per A
attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or gliagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opiold Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opiolds for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Dally Dose of Opioids for Safer Dos if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	Tes Yes	No No
attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the importance of I	attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Oploid Dosage Calculator and CDC handout "Calculating Total Dally Dose of Oploids for Safer Dos if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	Yes	No No
Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opiold Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opiolds for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dos if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	Yes	No
If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Dally Dose of Opioids for Safer Dos if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)		
7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Dally Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dos If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)		L
Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Dally Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dos If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	-	-
Some the reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Dally Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dos If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)		
Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dos If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	1	0
Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Oploid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Oploids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dos if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)		~40
Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Oploid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Oploids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dos if in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	IMPO	KIM
prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dos If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	_	0
Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dos If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation i section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	1	-
If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elemen of Good Faith Dispensing are met.)	If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation is section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)		
section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the element of Good Faith Dispensing are met.)	section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	age"	
(For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elemen of Good Faith Dispensing are met.)	For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.)	n note	es
에 하나요요 하다 가입니다. 이번 사람들은 사람들은 사람들은 이 사람들은 이 사람들은 이 사람들은 사람들은 사람들은 사람들이 되었다면 하나요요. 그는 사람들은 사람들은 사람들은 사람들은 사람들은 사람들은 사람들은 사람들은	of Good Faith Dispensing are met.)	1	
		e eiem	ents
	intes:	-	_
	19103.		
	Control of the Contro		
attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my	attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my		
professional judgment to review this prescription and I have:		0	

Proprietary & Confidential. This Checklist and the information contained herein is a Trade Secret of Walgreen Co.